



Paclitaxel protein bound (Abraxane®)

("pac-li-TAKS-el PROE-teen bownd")

How drug is given: by vein (IV)

Purpose: to kill cancer cells in breast cancer and other cancers

Things that may occur during or within hours of treatment

1. If you feel any burning or tingling near your IV, please tell your nurse right away. If you develop any swelling or redness after you go home, please call your doctor or nurse.
2. Your blood pressure may drop or your heart rate slow down while you are getting this drug. This usually goes back to normal on its own.
3. Some patients may have mild nausea. You may be given medicine to help with this.
4. Rarely, this drug may cause an allergic reaction. If you have a rash, facial flushing (warmth or redness of the face), or itching during or after you get this drug, please tell your doctor or nurse right away.

Things that may occur a few days to weeks later

1. You may lose some feeling, or have tingling or burning in your hands and/or feet. This is called peripheral neuropathy. This may increase with continued treatment. Please tell your doctor or nurse if you have trouble buttoning your clothes. Peripheral neuropathy should get better over time when the drug is stopped.
2. Loose stools or diarrhea may occur within a few days after the drug is started. You may take loperamide (Imodium A-D®) to help control diarrhea. You can buy this at most drug stores. Be sure to also drink more fluids (water, juice, sports drinks). If these do not help within 24 hours, **call your doctor or nurse.**
3. You may feel pain or weakness in your joints or muscles. This may happen about 2 to 3 days after you get paclitaxel protein bound (Abraxane®). This usually gets better in less than a week. If these bother you, ask your doctor or nurse what type of drugs you may use to help with this pain.

4. Your blood cell counts may drop. This is known as bone marrow suppression. This includes a decrease in your:
- Red blood cells, which carry oxygen in your body to help give you energy
 - White blood cells, which fight infection in your body
 - Platelets, which help clot the blood to stop bleeding

This may happen 7 to 14 days after the drug is given and then blood counts should return to normal. If you have a **fever of 100.5°F (39°C) or higher**, chills, a cough, or any bleeding problems, tell your doctor or nurse **right away**.

5. You may get mouth sores 7 to 10 days after this drug is given. It is important to keep your mouth clean. A soft-bristle toothbrush should be used for brushing your teeth. You may have a burning feeling and redness inside the mouth or on the lips. Use a **baking soda mouth rinse** 4 or 5 times a day.

Mouth Rinse Recipe:

Mix: 1 tsp **salt or baking soda** with 8 oz. glass of **water**

Other mouth rinses may be ordered by your doctor. Do not use mouth washes that have alcohol in them because they will dry out the mouth. If you cannot eat or swallow because of this, let your doctor or nurse know. Check with your doctor before having any dental work done.

6. Some or all of your hair may fall out around 3 to 4 weeks after treatment starts. You may lose hair on your head as well as facial and body hair. You may want to buy a wig before hair loss begins. Hair may grow back during treatment.
7. Some patients may feel very tired, also known as fatigue. You may need to rest or take naps more often. Mild to moderate exercise can also be helpful in maintaining your energy. Talk with your doctor or nurse.
8. You may have changes in your eyesight which include blurred vision. Eye exams may be done.
9. Skin changes, such as dryness or a rash on your body, may occur. Ask your doctor or nurse what lotions or creams you may use.
10. Your liver may be affected by this drug. Labs may be drawn to monitor your liver function.
11. You may have a fast or unusual heartbeat. If you feel any strange changes in your heartbeat, tell your doctor or nurse **right away**.

Things that may occur after your treatment ends (even months to years later)

Fertility:

- It is very important to use birth control if you are having sex since this drug could harm an unborn baby.
- Men may become sterile and may want to sperm bank prior to treatment.

- Women should not breastfeed while receiving this drug.
- For women, your period (menstrual cycle) may not be regular for a while or may stop completely and you may not be able to get pregnant.

If you have questions or concerns, please talk with your doctor or nurse.

The above information includes some, but not necessarily all, of the possible side effects of this medication. The side effects listed in this teaching sheet may not be the same ones you experience. Your side effects may be different, depending on how often you receive treatment (your schedule) and how much you receive each time (your dosage). Side effects may also vary if you are taking other medications. Please speak with your doctor or nurse if you have questions about possible side effects you may experience. This document should not take the place of conversations with members of your health care team.

If you experience any significant change in your health during or after treatment, contact a member of your health care team right away.

THIS SPACE RESERVED FOR WRITTEN COMMENTS OR NOTES FOR THE PATIENT AND FAMILY: